

# ATP1B1 Recombinant antibody

**Cat:**B36429S**Company:** HaoKebio**Uniprot ID:**P05026**Applications:** IHC:1:500-1:2000**Organism:**Rabbit

IHC-Polymer:1:2000-1:8000

**Species reactivity:**Human Mouse Rat

IHC-TSA:1:2500-1:10000

**Molecular Weight Calculation:** 35 kDa

IF:1:50-1:100

**Observed Molecular Weight:** 45-52 kDa

WB:1:2000-1:10000


**Background:**

ATP1B1 is one of beta subunits of the Na<sup>+</sup>/K<sup>+</sup> ATPase and responsible for formation and structural integrity of the Na<sup>+</sup>/K<sup>+</sup> ATPase. The Na<sup>+</sup>/K<sup>+</sup> ATPase is a plasma membrane pump consisting of alpha, beta, and gamma subunits. At least four of Na<sup>+</sup>/K<sup>+</sup>-ATPase beta subunits ( $\beta$ 1,  $\beta$ 2,  $\beta$ 3,  $\beta$ 4) have been identified in mammalian cells; the  $\beta$ 1-subunit (ATP1B1) is the most ubiquitous. The Na<sup>+</sup>/K<sup>+</sup> ATPase  $\beta$  subunits have multiple N-glycosylation sites. The predicted MW of ATP1B1 is 35 kDa, while it migrates around 40-52 kDa due to the variable glycosylation.

Store at -20 °C for one year.

**Experimental procedure:**

Antigen retrieval: Citrate buffer (pH 9.0) , Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

**Images:**

Sample: Mouse kidney, 4% PFA 12-24h

**Source of Reagents:**

发表[中文论文]请标注:ATP1B1(B36429S)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:ATP1B1(B36429S) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.

**Synonyms:**

242109D12, ATP1B 1, Beta 1 subunit of Na(+) K(+) ATPase, Na, K-ATPase  $\beta$ 1, Na<sup>+</sup>/K<sup>+</sup> ATPase beta 1 subunit

**Immunogen:**

Recombinant protein

**Isotype:**

IgG

**Subcellular location:**

Membrane,Cytoplasm

**Purity:**

Affinity purification

**Form:**

Liquid

**Storage Buffer:**

PBS with 0.02% sodium azide,100  $\mu$ g/ml BSA and 50% glycerol.

**Storage:**